



Chancroid - Reported cases: United States, 1981-1997

Cases (in thousands)



Ectopic pregnancy - Hospitalizations of women 15-44 years of age: United States, 1980-1996


Number of Ectopic Pregnancies



Note: Some variations in 1981 and 1988 numbers may be due to changes in sampling procedures. The relative standard error for these estimates ranges from 8% to 12%.

Source: National Hospital Discharge Survey (National Center for Health Statistics, CDC)

Genital herpes simplex virus infections - Initial visits to physicians' offices: United States, 1966-1997 and the Healthy People year 2000 objective




Source: National Disease and Therapeutic Index (IMS America, Ltd.)

Human papillomavirus (genital warts) - Initial visits to physicians' offices: United States, 1966-1997 and the Healthy People year 2000 objective



Source: National Disease and Therapeutic Index (IMS America, Ltd.)


Pelvic inflammatory disease - Initial visits to physicians' offices by women 15-44 years of age: United States, 1980-1997 and the Healthy People year 2000 objective



Source: National Disease and Therapeutic Index (IMS America, Ltd.)

Pelvic inflammatory disease - Hospitalizations of women 15-44 years of age: United States, 1980-1996

Hospitalizations (in thousands)




Note: The relative standard error for the estimates of the overall total number of PID cases range from 6% to 9%.

Source: National Hospital Discharge Survey (National Center for Health Statistics, CDC).

Trichomonal and other vaginal infections - Initial visits to physicians' offices: United States, 1966-1997


Visits (in thousands)



Source: National Disease and Therapeutic Index (IMS America, Ltd.)

Nonspecific urethritis - Initial visits to physicians' offices by men: United States, 1966-1997

Visits (in thousands)



Source: National Disease and Therapeutic Index (IMS America, Ltd.)